## **CLENPIQ Bowel Preparation**

	When	What to do before your procedure
	days before your procedure	<ul> <li>Pick up the prescription for CLENPIQ from your pharmacy.</li> <li>Do not eat nuts, seeds, whole wheat bread, beans, corn, raw fruits and vegetables with seeds or skin for 5 days before your procedure.</li> </ul>
1	day before your procedure	<ul> <li>Do not eat any solid food.</li> <li>If you have heart failure or kidney disease, follow your provider's instructions for your daily fluid amount.</li> <li>Only drink clear liquids, nothing red or purple. No alcohol.         <ul> <li>water</li></ul></li></ul>
	After midnight	<ul> <li>No dairy, coffee creamer, alternative dairy products such as oat milk, juice with pulp, chewing gum or candy.</li> <li>Step 1: Drink 1 bottle of CLENPIQ until it is empty. A straw may be helpful.</li> <li>Step 2: Drink five (5) 8 oz. cups of clear liquids within the next 5 hours.</li> <li>Stop eating any broth, popsicles, or JELL-O.</li> </ul>
	hours before your procedure	<ul> <li>Do not eat any solid food, broth, popsicles, or JELL-O.</li> <li>Drink the second bottle of CLENPIQ until it is empty.</li> <li>Then drink three (3) 8 oz. cups of clear liquids within the next 2 hours.</li> <li>Stool must be a clear, yellow-colored liquid (similar to apple juice) and without any solid material for your procedure.</li> <li>Take your approved medications with a small sip of water at least 3 hours before your procedure, unless otherwise directed by your provider.</li> <li>If you take diabetic medications, follow instructions given by your provider.</li> </ul>
	hours before your procedure	Stop drinking any liquids.

If you eat  $\underline{any}$  solid food on the day of your procedure, your procedure may be canceled.

If you have any questions, please contact the clinic.

